

NETTLE PAPER

**OUR GOAL
TO REVERSE NETTLE IN A CROP AVAILABLE FOR HUMAN IN ALL WORLD**

based on know-how

FROM TUBES THROUGH THE FIELD ON THE FORK

CONTENT

1.Opportunity	3
1.1 Burda´s farm and opportunity to use weedy nettle	3
1.2 Main benefits of nettle	3
2. Solution	6
2.1 Establishment of nettle field and testing food production	6
2.2 Clone preparation and reproduction	6
3.Our vision	9
4. References:	10

1.Opportunity

1.1 *Burda's farm and opportunity to use weedy nettle*

The farm cultivates on 4.5 ha agriculture soil in the status of ecological farming. Some from the fields are in the area where nettle (*Urtica dioica*, L.) weeding was a trouble. Due to the restriction of usage of the herbicides in ecological farming we were up for a challenge. How could these lands be beneficial? From agricultural historical data we found out that nettle cultivation is possible (for textile usage especially). For example, during World War I, both Germany and Austria were running out of cotton. In an effort to find a suitable substitute: the nettle. Moreover, nettle is a suitable plant as well as for ecological cultivation or permaculture (Vogl and Hartl, 2003). Many scientists have already tried to cultivate nettles. The main benefits are:

1. there is data on good yield without nitrogen fertilization (Kohler *et al.*, 1999).
2. the crop is based on several years (within 10-15 years it still achieves a satisfactory yield (Gatti *et al.*,2008)
3. green matter can be used in several ways (unsunked production to the food industry can be used in e.g. biogas, or used as a nutritionally full feed for livestock. This creates minimal losses as nettle is one of the low-cost plants)


Other advantages include:

- easy propagation and sustainable yields when using the principles of organic farming (Gatti *et al.*; 2008).
- as a perennial crop, it requires lower inputs in terms of soil processing, which maintains soil fertility and its structure (e.g. protects the top layer of soil against erosion, crusting and excretion (Szewczuk *et al.*2002).
- In addition to these effects, it also promotes biodiversity and acts as a food source for useful insects (e.g. water nymphs).
- is a fast-growing plant and therefore has an advantage over other weeds in the absorption of water and nutrients (Bacci *et al.*2009), so the use of chemicals against weeds is not necessary.
- the leaves are very rich in minerals and when they fall off, an important amount of nutrients are returned to the soil.

1.2 *Main benefits of nettle*


There are other reasons for growing nettle, it can be used whole - leaves, seeds, stem and root. In addition, many studies confirm very interesting compounds found mainly in nettle leaves, which have great potential for the food, pharmaceutical and cosmetic industries (tab. 1). Upton (2013) states that nettle has a high nutritional value and contains in its leaves: vitamins A, D and C, proteins, calcium

minerals, iron, potassium, manganese, choline, amines, antioxidant chlorophyll (picture 1) and 5-hydroxytryptophan.



Picture 1. Chemical scheme of chlorophyll. There are several types of chlorophyll, but all share the chlorin magnesium ligand which forms the right side of this diagram. Chlorophyll stimulates the immune system, or eliminates fungus in the body

Carotenoids (which include vitamin A) are a nutritionally important component, and a total of 9 have been identified in nettle leaves, the largest proportion being β -carotene (picture 2), which is a precursor of vitamin A.



Picture 2. Chemical scheme of β -carotene. It is a member of the carotenes, which are terpenoids (isoprenoids), synthesized biochemically from eight isoprene units and thus having 40 carbons. Among the carotenes, β -carotene is distinguished by having beta-rings at both ends of the molecule.

Guil-Guerrero *et al.*, 2003 states that the total amount of carotenoids in fresh leaves is 29.6 mg 100⁻¹ g dry weight. As is known their fundamental contribution lies in antioxidant action. This means that it protects cells from damage from unstable oxygen molecules, so-called free radicals. Rafajlovska *et al.* (2001) states that extracts of nettle plants contain 6.8% palmitic, 1.1% stearic, 3.6% oleic, 20.2% linoleic and 12.4% linoleic acid. These fatty acids are an important component of lipids, and some of them are essential. They are important sources of energy for humans and animals too. Therefore, when they are metabolized, they yield large quantities of ATP.

A lot of studies have shown that *Urtica dioica* can be used in medicine. The aqueous and alcoholic extracts have been used for hundreds of years for the treatment of anaemia (Pinelli *et al.*, 2008), rheumatism (Jaric' *et al.*, 2007), gout and eczema (Orčic' *et al.*, 2014), and treatment of urinary, bladder and kidney problems (Orčic' *et al.*, 2014). Beneficial effects have also been reported on inflammation, hypoglycaemia, hypotension, benign prostatic hyperplasia, cardiovascular problems, arthritis, and allergic rhinitis (Upton, 2013). Furthermore, stinging nettle exhibits antioxidant, antimicrobial, antifungal, antiviral, and antiulcer activity (Upton, 2013). Nettles possess noticeable antimicrobial activity against Gram-positive and Gram-negative bacteria when compared with standard and strong antimicrobial compounds, such as miconazole nitrate, amoxicillin-clavulanic acid, ofloxacin and netilmicin (Gülçin *et al.*, 2004). Different fractions of various *Urtica* species have been studied to determine their antimicrobial activity. The results indicate the great potential of this plant for the discovery of novel effective compounds.

Tab. 1 Chemical composition of nettle leaf powders (Adhikari *et al.* 2016)

Parameter	Content
Moisture (%)	7.04±0.77
Crude protein (%)	33.77±0.35
Crude fiber (%)	9.08±0.14
Crude fat (%)	3.55±0.06
Total ash (%)	16.21±0.54
Carbohydrate (%)	37.39±0.72
Calcium (mg/100 g)	168.77±1.47
Iron (mg/100 g)	227.89±0.21
Tannins (%)	0.93±0.01
Polyphenols (mg GAE/g)	128.75±0.21
Carotenoids (µg/g, db)	3496.67±0.56
Caloric value (kcal/100 g)	307.24±0.13

Nettle provides humans and animals with nutrients and bioactive components, which support antimicrobial activity, immune enhancement and stress reduction. It is worth noting that the World Health Organization (WHO), in its monographs on 'Selected medicinal plants', describes *Urticae* as valuable herbs for many medicinal uses. The European Commission Directorate-General For Health and Food Safety showed that *Urtica* spp. fulfils the criteria of a foodstuff, as defined in Regulation (EC) No. 178/2002. This opinion is supported by the European Food Safety Authority (EFSA). It concluded that nettle has neither an immediate nor delayed harmful effect on human or animal health and has no negative effect on the environment.

2. Solution

2.1 Establishment of nettle field and testing food production


We established a testing cultivation nettle field. The first yield was about 9tha^{-1} . There was information about cultivation of stinging nettle for the textile industry during the second war in Germany. In 1940 stinging nettle was joined to the wild plant breeding programme for fibre production at the Institute of Applied Botany in Hamburg, which managed to derive several clones with a high fiber content and in this point of view, that could be considered an agricultural crop for this purpose. These clones are still kept in German and Austrian research institutions (Vogl and Hartl, 2003). At the start, we took this historical agrotechnical data and we tried to apply them on our conditions. However, the chemical composition of plants is affected by different factors, including the variety, genotype, climate, soil, vegetative stage of the plant, harvest time, storage, processing and treatment (Angela and Meireles, 2009). When and how nettles are harvested strongly determines the final product. For example, for fiber production, stinging nettles should be harvested when the seeds are mature or when the stalks reach 80% of the aboveground biomass, from the second year of planting. During the first year, the stalks are too thin, too ramified and have too many leaves. If the main product is to be the leaves, younger plants are harvested. The time of year for nettle harvesting depends on the purpose. Plants collected in April are used for fodder, medicine or chlorophyll production. Nettles harvested at the end of June are used for fiber production. The second harvest in September may be used for the collection of leaves (Di Virgilio *et al.*, 2015). It seems that quality and chemical composition of stinging nettle will be affected by epigenetic factors. It means we are able by suitable chosen agrotechnology to produce suitable plant material for food production in this way we can influence the taste of final products (eg. hydroponic or aquaponic culture, as well as indoor technology).

It is important to realize that a big advantage of nettle cultivation is that harvest could be used for plenty of usage: feed, fertilizer, mass into biogas, textile and last not least food. Therefore, nettle is appropriate for a circular economy (more detail in business plan).

After harvesting non-cultivated nettle (heterogeneous material), we started to test if it would be possible to make food products (pasta, lemonade etc.) based on different conditions of agrotechnology. After we fine tuned agrotechnology and food technology, we found the taste, which is interesting and moreover the nettle belongs to a **superfood** (Kriegel *et al.*, 2018). **Our nettle products perfectly fit a healthy lifestyle.** Our pilot food products we released in the market and customers reacted in a positive way. Based on this data we did a market research and we revealed a gap - there are a few food products from nettle and all of them are not customer convenience (more details in a section Business plan). Therefore we take the advantage. We found out we need to have nettle as a crop.

2.2 Clone preparation and reproduction

However, we did not have homogenous material due to the dioecy of nettle, which has a separate male and female sex. Closely, Stinging nettle (picture 3) is a perennial herbaceous plant belonging to the Urticaceae family.



Picture 3. Plant habitus of *Urtica dioica*

It is a well-known and common species, spread in temperate and tropical zones of Europe, Asia and America, adapted to a variety of climatic conditions. Stinging nettle is a perennial, monoecious plant, flowering and fruiting in summertime. Its stems and leaves are covered by stinging trichomes containing a fluid which causes blistering when entering the skin (Bisht *et al.*, 2012). Surprisingly, there are a number of subspecies for this species, probably due to polyploidization; most plants are predominantly tetraploid in nature (Rejlová *et al.*, 2019). Genetic analyses point to a model of gender inheritance that is not bound to sex chromosomes and is predominantly Mendelistically controlled by one gene, however, there are exceptions to this model (Glawe and Jong, 2008). For this reason, homogenisation of nettle is difficult, however by using in vitro method, it is possible. In vitro plant tissue culture is a collection of techniques used to maintain or grow plant cells, tissues or organs under sterile conditions on a nutrient culture medium of known composition. It is widely used to produce clones of a plant in a method known as micropropagation. Plant tissue culture is used widely in the plant sciences, forestry, and in horticulture. for nettle Applications we can use:

- **the commercial production of plants used as potting, landscape, and florist subjects, which uses meristem and shoot culture to produce large numbers of identical individuals -clones - (already done)**
- to conserve rare or endangered nettle species
- a plant breeder may use tissue culture to screen cells rather than plants for advantageous characters, e.g. salt resistance/tolerance
- large-scale growth of plant cells in liquid culture in bioreactors for production of valuable compounds, like plant-derived secondary metabolites and recombinant proteins used as biopharmaceuticals
- to cross distantly related species by protoplast fusion and regeneration of the novel hybrid

- to rapidly study the molecular basis for physiological, biochemical, and reproductive mechanisms in plants, for example in vitro selection for stress tolerant plants
- for chromosome doubling and induction of polyploidy, for example doubled haploids, tetraploids, and other forms of polyploids. This is usually achieved by application of antimetabolic agents such as colchicine or oryzalin.
- as a tissue for transformation, followed by either short-term testing of genetic constructs or regeneration of transgenic plants

Therefore we started a biotechnological approach in collaboration with Mendel university in Brno. Based on in vitro technology, our coopartner was able to generate genetic homogenous nettle called clones (picture 4). Because we have obtained knowledge about nettle habitus connected to food properties from field observation, we could provide appropriate candidates for micropropagation.



Picture 4. Nettle clones growing in vitro culture (Ing. Kamila Lónová, Department of Plant Biology, Mendel university in Brno)

After acclimatization we obtained special clones to make our growth homogenous. Nowadays, we cultivate these clones in our field Vlčice (picture 5). We are able to trade these clones thanks to nettle nursery and as well as to offer consultancy regarding the nettle cultivation.



Picture 5. Nettle field in the Vlčice area

Our main know-how is the production of 3 products: juice - spinach - cold syrup suitable for retail packaging as well. Our production process is specific in 3 ways of fermentation. We are able to preserve salt, fat, acid and freeze. Moreover, we can produce as well as nettle extract for cosmetics and pharmacy. Furthermore, we processed the protocol for nettle explant, so we have opened the door for the breeding program of nettle by using a biotechnological approach, which makes our expansion easier.

3. Our vision

Our potencial, we see in the combination of new disciplines of biotechnology, nanotechnology and blockchain. Our vision is to make quality nettle products which will be fitted for food, textil, cosmetics and pharmaceutical ware based on knowledge from genomics or nanotechnology and their tools. To realize our vision we need an economic tool making external markets available, therefore we take advantage of blockchain and token economy.

BIOTECHNOLOGY, NANOTECHNOLOGY AND OUR VISION (FROM TUBES THROUGH THE FIELDS ON THE FORKS AND ON BODY)

Clones are a future for nettle farming due to obstacles with homogenization due to dioecy of *Urtica dioica*. Our plan is to do deeper genomic analysis based on next generation sequencing. We have already processed genome sequencing by an Illumina method (picture 6) to see basic differences between male and female (the cooperation with The Application Laboratory for Agricultural Research). Based on this data, we clearly see that transcriptome sequencing will be needed to reveal candidate genes playing a crucial role in metabolic pathways which are targeted. After identification of these genes we will be able to produce plants with novel and extraordinary traits.

Target metabolic pathways for genetic engineering:

- phenolic pathways
- proteosynthesis
- metabolisms of flavonoids, tannins
- fatty acids synthesis
- metabolic pathways in trichomes

We would like to be a crucial player in the nettle products delivery in the world. To realize our vision we need an economic tool making external markets available, therefore we take advantage of blockchain and token economy. Among nettle food products you will find such products that even classic large food businesses or food retail companies need not be ashamed of. They are both conventional and organic products. Superfood, kitchen ingredients and semi-finished products, products for direct consumption originating from ecologically clean locations, of course gluten-free and non-cacate products, vegetarian and vegan foods. Typical products of bakers, confectioners, liqueurs, brewers, milkmen, cheesemakers, butchers, smokers. Not only these nominal products can find an excellent basis or ingredient in the technologically well-processed and prepared nettle.

4. References:

Adhikari, B.M.; Bajracharya, A.; Shrestha, A.K. 2016. Comparison of nutritional properties of stinging nettle (*Urtica dioica*) flour with wheat and barley flours. *Food Sci. Nutr.* 4, 119–124.

Angela, M.; Meireles, A. *Extracting Bioactive Compounds for Food Products: Theory and Applications*; CRC Press: Boca Raton, FL, USA, 2009.

Bacci, L., Baronti, S., Predieri, S., di Virgilio, N., 2009. Fiber yield and quality of fiber nettle (*Urtica dioica* L.) cultivated in Italy. *Ind. Crops Prod.* 29, 480–484.

Di Virgilio, N.; Papazoglou, E.G.; Jankauskiene, Z.; Di Lonardo, S.; Pralczyk, M.; Wilegusz, K. 2015. The potential of stinging nettle (*Urtica dioica* L.) as a crop with multiple uses. *Ind. Crops Prod.* 68, 42–49

Filipek, T., Badora, A., Kaczor, A., Krawiec, Z., 1999. *Podstawy i skutki chemizacji agroekosystemów*. Wydawnictwo Akademii Rolniczej, Lublin

Gatti, E., Di Virgilio, N., Baronti, S., Bacci, L., 2008. Development of *Urtica dioica* L. propagation methods for organic production of fiber. In: *Proceedings of the 16th IFOAM Organic World Congress*, Modena, June 16–20, pp. 112–113.

Glawe, G.A., de Jong, T.J., 2009 Complex sex determination in the stinging nettle *Urtica dioica*. *Evol Ecol* 23, 635.

Guil-Guerrero, J.L.; Reboloso-Fuentes, M.M.; Isasa, M.E.T. 2003 Fatty acids and carotenoids from stinging nettle (*Urtica dioica* L.). *J. Food. Compos. Anal.* 16, 111–119.

Gülçin, I.; Küfrevioğlu, O.I.; Oktay, M.; Büyükkuroğlu, M.E. 2004. Antioxidant, antimicrobial, antiulcer and analgesic activities of nettle (*Urtica dioica* L.). J. Ethnopharmacol. 90, 205–215.

Gülçin, I.; Küfrevioğlu, O.I.; Oktay, M.; Büyükkuroğlu, M.E. 2004. Antioxidant, antimicrobial, antiulcer and analgesic activities of nettle (*Urtica dioica* L.). J. Ethnopharmacol. 90, 205–215..

Jarić, S., Popović, Z., Macukanović-Jović, M., Djurdjević, L., Mijatović, M., Karadžić, B., Mitrović, M., Pavlović, P., 2007. An ethnobotanical study on the usage of wild medicinal herbs from Kopaonik Mountain (Central Serbia). J. Ethnopharmacol. 111, 160–175.

Köhler, K., Schmidtke, K., Rauber, R., 1999. Eignung verschiedener Pflanzenarten zur Untersaat in Fasernesseln (*Urtica dioica* L.). In: Hoffmann, H., Müller, S. (Eds.), Beiträge Zur 5. Wissenschaftstagung Zum Ökologischen Landbau. Koster Verlag, Berlin, Germany, pp. 496–500.

Kregiel D, Pawlikowska E, Antolak H, 2018 *Urtica* spp.: Ordinary Plants with Extraordinary Properties. *Molecules*. 23(7):1664.

Orčić, D.; Francišković, M.; Bekvalac, K.; Svirčev, E.; Beara, I.; Lesjak, M.; Mimica-Dukić, N. 2014 Quantitative determination of plant phenolics in *Urtica dioica* extracts by high-performance liquid chromatography coupled with tandem mass spectrometric detection. Food Chem. 143, 48–53.

Pinelli, P.; Ieri, F.; Vignolini, P.; Bacci, L.; Baroni, S.; Romani, A., 2008 Extraction and HPLC analysis of phenolic compounds in leaves, stalks, and textile fibers of *Urtica dioica* L. J Agric. Food Chem. 56, 9127–9132.

Rafajlovska, V.; Djarmati, Z.; Najdenova, V.; Cvetkov, L. Extraction of stinging nettle (*Urtica dioica* L.) with supercritical carbon dioxide. Presented at the 1st Black Sea Basin Conference on Analytical Chemistry, Odessa, Ukraine, 11–15 September 2001.

Rejlová L., Chrtěk J., Trávníček P, Lučanová M, Vít P., Urfus T. Polyploid evolution: The ultimate way to grasp the nettle. 2019. PLoS One 14, 24, doi: 10.1371/journal.pone.0218389

Szewczuk, C., Stepniak, M., Sugier, D., 2002. Zawartość wybranych związków organicznych i mineralnych w części nadziemnej pokrzywy zwyczajnej (*Urtica dioica* L.) w zależności od fazy rozwojowej zbieranych roślin. Acta Sci. Pol. Agric. 1, 163–169.

Upton, R., 2013. Stinging nettles leaf (*Urtica dioica* L.): extraordinary vegetable medicine. J. Herb. Med. 3, 9–38.

Vogl, C.R., Hartl, A., 2003. Production and processing of organically grown fiber nettle (*Urtica dioica* L.) and its potential use in the natural textile industry: a review. Am. J. Altern. Agric. 18, 119–128.